

Your surgery is scheduled at Columbia Eye Surgery Center, 1920 Pickens Street, Columbia, S.C. 29201

On: \_\_\_\_\_ Please arrive at \_\_\_\_\_. You will be called with your time for arrival one to three days prior to surgery. Family members should be prepared to wait about three hours for surgery patients.

CHECKLIST

1. NOTHING TO EAT OR DRINK AFTER MIDNIGHT THE NIGHT PRIOR TO SURGERY.

2. Take your regular medications the morning of surgery with a SMALL SIP OF WATER > EXCEPT \_\_\_\_\_

\*\*\*\*\*

\*\*\*\*\* DIABETIC PATIENTS \*\*\*\*\*

DO NOT TAKE YOUR INSULIN OR DIABETIC TABLET ON THE DAY OF SURGERY UNLESS OTHERWISE INSTRUCTED.

\*\*\*\*\*

3. Bring all BLOOD PRESSURE and /or DIABETIC medications (including your INSULIN ) with you to the Surgery Center in their originally labeled containers the day of surgery.

4. Begin using Besivance (or Moxifloxacin or Ofloxacin )- 3 times a day AND Prolensa (or Ilevro ) 1 time

a day in your \_\_\_\_\_ eye beginning on Saturday \_\_\_\_\_, 3 days before surgery.

\*\*\*WAIT AT LEAST 5 MINUTES BETWEEN DROPS\*\*\*

5. Wear loose fitting, comfortable clothing. Please wear a short sleeve top that buttons up the front without a T-shirt. Leave all valuables at home.

6. Wash your face with just soap and water. DO NOT wear make-up, lotions or creams the morning of surgery.

7. Bring a responsible adult to drive you home after surgery. You will also need someone to bring you back to the office the next day.

8. You will be given a kit to take home with you the day of surgery. Please bring the kit and your drops with you the next day so we can give you post-operative care instructions.

9. You will start using the Prednisolone drops 3 times a day into the operative eye starting after surgery.